

The role of interleukin 4 and IL-4RA in intervertebral disc degeneration: investigation of single nucleotide polymorphisms in genes and a systematic review & meta-analysis of IL-4 expression level

Sara Hanaei^{a,b,c}, Sina Abdollahzade^d, Maryam Sadr^a, Mohammad Hossein Mirbolouk^e, Ehsan Fattahi^e, Alireza Khoshnevisan^e, Nima Rezaei^{b,c,f}

^aMolecular Immunology Research Center, Tehran University of Medical Sciences, Tehran, Iran

^bResearch Center for Immunodeficiencies, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

^cNetwork of Immunity in Infection, Malignancy and Autoimmunity (NIIMA), Universal Scientific Education and Research Network (USERN), Tehran, Iran

^dDivision of Neurosurgery, Department of Surgery, Rajayi Hospital, Qazvin University of Medical Sciences, Qazvin, Iran

^eDepartment of Neurosurgery, Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran

^fDepartment of Immunology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Corresponding author: Nima Rezaei, Research Center for Immunodeficiencies, Children's Medical Center, Tehran University of Medical Sciences, Dr Qarib St, Keshavarz Blvd, Tehran, Iran; email: rezaei_nima@tums.ac.ir

Abstract

Background: Intervertebral disc degeneration (IVDD) is a multifactorial disease that is sensitive to the balance between anti-inflammatory and pro-inflammatory cytokines. This study investigated the single nucleotide polymorphisms (SNPs) of interleukin 4 (IL-4) in IVDD. **Methods:** Genomic DNA of peripheral mononuclear cells of 76 IVDD patients and 140 healthy controls were investigated for three SNPs of IL-4 (rs2243248 (-1098G/T), rs2243250 (-590 C/T), rs2070874 (-33 C/T)) and 1 SNP of IL-4RA (rs180275, +1902 A/G) through PCR-SSP method. **Results:** The 'C' allele frequency of IL-4 rs2243250 was 104 in 76 patients, while it was 149 in 140 controls (OR = 2, p = .001); also this SNP was significantly associated with post-operative pain reduction. The 'C' allele of IL-4 rs2070874 (130 in 76 patients, and 200 in 140 controls, OR = 2.66), and the 'CC' genotype were more frequent among patients (OR = 3.98, p < .001) than controls. 'TTT' haplotype was more common in controls (OR = 0.36, p < .001) and 'TCC' was also more common in patients (OR = 1.75, p = .012). A meta-analysis of previous studies found significantly higher IL-4 levels in disc tissues of IVDD patients, which was not similarly found in blood samples. **Conclusion:** The immune system plays an important role in IVDD. The extent and progress of the disease vary significantly with IL-4 level. Meanwhile, the rs2070874 and rs2243250 SNPs of IL-4 were significantly associated with IVDD in Iranian patients.

Keywords: Intervertebral disc degeneration, single nucleotide polymorphism, interleukin 4, Interleukin-4 receptor alpha, immunogenetics, cytokine